

ON AMBIENT EMBEDDINGS IN THE HILBERT CUBE

David G. WRIGHT

Department of Mathematics, Brigham Young University, Provo, UT 84602, USA

Received 12 March 1985

It is shown that for each closed subset X of codimension at least three in the Hilbert cube Q there is a Cantor set C in Q so that no homeomorphism of Q onto itself takes C into X .

AMS(MOS) Subj. Class.: Primary 54C25, 57N35;
Secondary 57N20

ambiently universal set	Cantor set
ambient embedding	Hilbert cube

1. Introduction

Let M and N be subsets of a topological space X . We say M is *ambiently embeddable* in N if there is a homeomorphism h of X onto itself that carries M into N . We call a subset U of X an *ambiently universal set* for a family F if each set in F is ambiently embeddable in U .

Bothe has made an extensive study of ambiently universal sets in E^n [2, 3, 4, 5]. Recently the author [13] showed that each closed set X in E^n of dimension at most $n-3$ fails to be ambiently universal with respect to Cantor sets in E^n . From these works our understanding of ambiently universal sets in E^n is fairly complete.

The purpose of this note is to generalize the results of [13] to the Hilbert cube.

2. Definitions and notation

We let S^n , B^n , and E^n denote the n -sphere, the n -ball, and Euclidean n -space, respectively. We let I denote the interval $[-1, 1]$. The product of n copies of I will be denoted by I^n , and the Hilbert cube which is the product of countably infinite copies of I will be denoted by Q . A closed subset X of Q is said to have *codimension* $\geq k$ if $H_q(U, U-X) = 0$ for $0 \leq q < k$ and all open subsets U of Q , the

homology being taken with integer coefficients. We say X has *codimension* k if X has $\text{codimension} \geq k$ but does not have $\text{codimension} \geq k+1$ [9, 11].

3. Bing Cantor sets in Q

In E^n ($n \geq 3$) wild Cantor sets may be described as the intersection of nested compact manifolds M_i so that each component of each M_i is homeomorphic to $B^2 \times S^{n-2}$ and contains exactly two components of M_{i+1} [1, 12].

Daverman has given a clever way of generalizing the Bing Cantor sets to the Hilbert cube [8]. Consider Bing's decomposition G of the 3-cube I^3 into points and tame arcs [1]. Bing showed that the decomposition space I^3/G is homeomorphic to an Alexander crumpled cube and then showed that two copies sewn together with the identity homeomorphism along the boundary yields a 3-sphere. Daverman considered the decomposition G' of $I^3 \times Q$ consisting of points and (arcs of G) $\times Q$. He showed that $(I^3 \times Q)/G'$ is homeomorphic to Q . The image of the nondegenerate elements of G' in Q forms a Cantor set that we also call a *Bing Cantor set*.

4. Ramified Bing Cantor sets in Q

By a *3-ball with end disks* we mean a 3-ball B that has two disjoint disks D_1 and D_2 identified in the boundary of B . A 3-ball with end disks may be identified with $B^2 \times I$ in such a manner that the end disks are identified with $B^2 \times \{-1\}$ and $B^2 \times \{1\}$. Whenever a 3-ball is given in a factored form $D^2 \times [a, b]$, where D^2 is a 2-dimensional disk and $[a, b]$ is a closed interval, we consider $D^2 \times [a, b]$ to be a 3-ball with end disks $D^2 \times \{a\}$ and $D^2 \times \{b\}$.

Consider the 3-ball $B^2 \times I$. Consider n disjoint disks D_1, D_2, \dots, D_n in the interior of B^2 . We call $\bigcup D_i \times I$ an n -fold ramification of $B^2 \times I$ [8, 10, 13].

Consider the 3-balls T_1 and T_2 in $B^2 \times I$ as shown in Fig. 1. The end disks of T_1 and T_2 are the intersections of T_1 and T_2 with the boundary of $B^2 \times I$. We call the embedding of $T_1 \cup T_2$ in $B^2 \times I$ a Bing embedding.

Let D^2 be a disk in the interior of I^2 and set $W_0 = D^2 \times I \subset I^3$. Inductively define W_{i+1} to be a finite collection of 3-balls with end disks by taking a Bing embedding in each component of W_i . The Bing decomposition of I^3 can be described as points and the components of $\bigcap W_i$. If care is taken the components of $\bigcap W_i$ will be arcs. The arcs simplify our description but are not essential to our proof.

Let $\alpha = a_0, a_1, a_2, \dots$ be a sequence of positive integers. We construct a ramified Bing decomposition with respect to α of I^3 as points and the components of the intersection of nested manifolds M_0, M_1, M_2, \dots . The set M_0 is, as before, $D^2 \times I$. Each component of each M_i is a 3-ball with end disks. Let i be a nonnegative integer. The manifold M_{2i+1} is obtained by taking an a_i -fold ramification in each




Fig. 1.

component of M_{2i} . The manifold M_{2i+2} is obtained by taking a Bing embedding in each component of M_{2i+1} . If care is taken each component of $\bigcap M_i$ will be an arc.

Let G be a ramified Bing decomposition of I^3 . The Daverman argument also shows that the decomposition G' of $I^3 \times Q$ consisting of points and $(\text{arcs of } G) \times Q$ yields a space homeomorphic to Q . Once again the image of the nondegenerate elements is a wild Cantor set which we call a ramified Bing Cantor set.

5. Grotes and Bing decompositions

An n -stage grope is a 2-dimensional polyhedron A_n along with a decomposition into subpolyhedra $A_{-1} \subset A_0 \subset A_1 \subset A_2 \subset \dots \subset A_n$ so that A_0 is a disk with handles whose boundary is A_{-1} . For $0 \leq i \leq n$, $A_i - A_{i-1}$ is a finite collection of open disks with handles. The closure of each component of $A_i - A_{i-1}$ is a closed disk with handles whose intersection with A_{i-1} is a handle curve of $A_{i-1} - A_{i-2}$ for $i > 0$, the correspondence between components of $A_i - A_{i-1}$ and handle curves of $A_{i-1} - A_{i-2}$ being one-to-one. See [6] for more details.

Theorem 5.1. *Let M_i be a defining sequence for a ramified Bing decomposition of I^3 corresponding to the sequence $\alpha = a_0, a_1, a_2, \dots$. Let A_n be an n -stage grope with decomposition A_i . If $f: A_n \rightarrow I^3 - \bigcap M_i$ is a continuous function so that $f(A_{-1}) \subset I^3 - M_0$ and $f|_{A_{-1}}$ is essential in $I^3 - M_0$, then $a_n \leq$ twice the number of handles of $A_n - A_{n-1}$.*

The proof of Theorem 5.1 is implicit in [13]; however, some comment is in order. The proofs in [13] avoid $n = 3$. These proofs deal with Bing decompositions of E^n where the nondegenerate elements are described by manifolds of the form $B^2 \times S^{n-2}$.

For $n > 3$ the fundamental group of the boundary of $B^2 \times S^{n-2}$ is isomorphic with the integers. For $n = 3$ this is not the case. However, appropriate modifications do work for the Bing decomposition of I^3 instead of E^3 .

Perhaps the easiest way to verify Theorem 5.1 is to take the 2-spin [7] of the ramified Bing decomposition of I^3 . This gives a decomposition of S^5 that is of the type studied in [13] and the function f can be approximated by an embedding. Then Theorems 7.5, 7.8, and 8.2 of [13] may be invoked directly.

6. The main theorem

Theorem 6.1. *Let X be a closed subset of Q that has codimension ≥ 3 . There is a Cantor set C in Q so that C is not ambiently embeddable in X .*

Proof. Let J_0, J_1, J_2, \dots be a collection of simple closed curves in $Q - X$ that are dense in the space of embeddings of S^1 into Q . For each nonnegative integer n , let P_n be an n -stage grope in $Q - X$ that has a decomposition $A_{-1} \subset A_1 \subset \dots \subset A_n = P_n$ so that $A_{-1} = J_n$. The J_n and P_n exist because X has codimension ≥ 3 . Let b_n be the number of handles of $A_n - A_{n-1}$ and set $a_n = 2b_n + 1$.

Let M_i be a defining sequence for a ramified Bing decomposition of I^3 with respect to the sequence a_0, a_1, a_2, \dots .

Let G be the resulting decomposition of I^3 and G' the decomposition of $I^3 \times Q$ described by Daverman. We identify $(I^3 \times Q)/G'$ with Q . The image of the nondegenerate elements of G' is a ramified Bing Cantor set C in Q . Suppose C can be pushed into X by a homeomorphism of Q . Then we may assume that the identification of $(I^3 \times Q)/G'$ with Q takes the image of the nondegenerate elements of G' into X . Hence, with this identification, we may consider the sets J_i and P_i to be contained in $I^3 \times Q$ and missing the nondegenerate elements of G' . Since the simple closed curves J_i are dense, for some n the projection $\pi: I^3 \times Q \rightarrow I^3$ restricted to P_n gives a map of a grope into $I^3 - \bigcap M_i$ so that $\pi(J_n) \cap M_0$ is empty and $\pi|_{J_n}$ is essential in $I^3 - M_0$. Theorem 5.1 implies that $a_n \leq 2b_n$. But $a_n = 2b_n + 1$, a contradiction. \square

It is known [14] that a closed set in Q that is not strongly infinite dimensional must have infinite codimension. Hence we get the following.

Corollary 6.2. *Let X be a closed subset of Q that is not strongly infinite dimensional (this includes finite dimensional and countably infinite dimensional sets). Then there is a ramified Bing Cantor set C so that C is not ambiently embeddable in X .*

References

- [1] R.H. Bing, A homeomorphism between the 3-sphere and the sum of two solid horned spheres, *Ann. of Math.* 56 (2) (1952) 354–362.

- [2] H.G. Bothe, Universalmengen bezüglich der Lage im E^n , *Fund. Math.* 56 (1964) 203–212.
- [3] H.G. Bothe, Zur Lage null- und eindimensionaler Punktmengen in E^3 , *Fund. Math.* 58 (1966) 1–30.
- [4] H.G. Bothe, Zwei Universalmengen in euklidischen Raum, *Math. Nachr.* 39 (1969) 117–133.
- [5] H.G. Bothe, Ein eindimensionales Kompaktum in E^3 , das sich nicht lagetreu in die Mengersche Universalkurve einbetten lässt, *Fund. Math.* 54 (1964) 251–258.
- [6] J.W. Cannon, The recognition problem: What is a topological manifold?, *Bull. Amer. Math. Soc.* 84 (1978) 832–866.
- [7] J.W. Cannon and R.J. Daverman, Cell-like decompositions arising from mismatched sewings: applications to 4-manifolds, *Fund. Math.* 111 (1981) 211–233.
- [8] R.J. Daverman, Infinite inflations of crumpled cubes, *Topology and its Applications* 12 (1981) 35–43.
- [9] R.J. Daverman and J.J. Walsh, Čech homology characterizations of infinite dimensional manifolds, *Amer. J. of Math.* 103 (1981) 411–435.
- [10] W.T. Eaton, A generalization of the dog bone space to E^n , *Proc. Amer. Math. Soc.* 39 (1973) 379–387.
- [11] D.J. Garity, Aleksandrov's problem, Pol's example and an alternate approach to dimension, *Proceedings of the BYU Topology Conference 1984*, to appear.
- [12] L.L. Lininger, Actions on S^n , *Topology* 9 (1970) 301–308.
- [13] D.G. Wright, Ambiently universal sets in E^n , *Trans. Am. Math. Soc.* 277 (1983) 655–664.
- [14] D.G. Wright, On sets with finite codimension in the Hilbert cube, *Proceedings of the BYU Topology conference 1984*, 17–23.